

Is dietary pesticide exposure related to semen quality? Positive evidence from men attending a fertility clinic

Hagai Levine^{1,2,*} and Shanna H. Swan²

¹Braun School of Public Health and Community Medicine, Hebrew University-Hadassah and The Hebrew University Center of Excellence in Agriculture and Environmental Health, Jerusalem, Israel ²Department of Preventive Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

*Correspondence address. Department of Preventive Medicine, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1057, New York, NY 10029, USA. Tel: +1-212-824-7184; Fax: +1-212-996-0407; E-mail: hagai.levine@mssm.edu

Submitted on February 25, 2015; resubmitted on February 25, 2015; accepted on February 28, 2015

Chiu and her colleagues from Harvard School of Public Health and Massachusetts General Hospital present important new data on semen quality in relation to dietary pesticide exposure via fruit and vegetable intake (Chiu *et al.*, 2015). To examine this relationship the authors utilized a novel approach that classifies fruits and vegetables into high versus low-to-moderate pesticide residue groups based on data from the United States Department of Agriculture (USDA) Pesticide Data Program (PDP). Data obtained from a validated food frequency questionnaire (FFQ) combined with USDA PDP yielded individual measures of intake of fruits and vegetables with higher pesticide residues. The authors found that semen quality was reduced among men in the highest quartile of exposure, a finding which could have clinical and public health implications.

Pesticides are designed to be biocides and an extensive literature demonstrates that many of these chemicals adversely impact human reproductive function. It has long been known that occupational exposure to certain pesticides have a devastating effect on semen quality and male fertility. For example plantation workers exposed to the nematocide 1,2-dibromo-3-chloropropane (DBCP) were rendered azoospermic (Whorton *et al.*, 1977). However, studying the subtler effects of non-occupational pesticide exposure on semen quality remains a challenge that is addressed by Chiu and colleagues using a novel method of exposure assessment. Their compelling results justify a discussion of methodological issues and public health implications.

Methodological issues

Interest in the association between pesticide exposure and semen quality is growing rapidly and over 1000 publications in this field have been published during the last 50 years (Fig. 1). This is a diverse literature, heterogeneous in populations, exposures, methods and study results. In light of this literature, we discuss briefly important methodological issues in this study: exposure assessment and timing, population selection, semen analysis methods, sample size and statistical analysis.

The authors have tackled the challenge of estimating exposure by coupling USDA PDP surveillance data with FFQ-based intake of fruits and vegetables. This method has been used to examine associations with other outcomes, but not, to our knowledge, semen parameters. For example, Curl *et al.* recently showed an association between urinary dialkyl phosphate concentrations and estimated dietary organophosphate exposure based on FFQ and USDA PDP pesticide measures (Curl *et al.*, 2015). In a systematic review of environmental and occupational pesticide exposure and semen quality, 15 of the 17 studies included used biomonitoring, while the others used questionnaire data and work histories (Martens and Perry, 2013). While exposure assessment by human biomonitoring may be the gold standard, high costs, the need to collect and store urine samples and the requirement of a qualified lab limit its utility. Moreover, high analytic costs preclude analysis of multiple samples, and basing exposure assessment on a single spot sample has its own limitation. Assessment of pesticide exposure with the aid of a pesticide residue surveillance program is an alternative method, and the one employed by Chiu and colleagues. This strategy could be especially useful in countries with high exposure to pesticides through diet, such as in Israel, where associations were found between fruit consumption and urinary organophosphate pesticide metabolite concentrations using biomonitoring (Berman *et al.*, 2013).

Timing of exposure is an important consideration in sperm studies, as exposure effects may differ during spermatogenesis (70–75 days). In Chiu *et al.* men completed their FFQ up to 18 months prior to semen collection. The authors examined the question of timing of exposure and found that results were similar in the full cohort and when restricting analysis to men whose semen sample was collected within 1 year of completing the FFQ. Therefore, in this analysis exposure misclassification due to improper timing is unlikely.

Chiu and colleagues recruited male partners from couples presenting to a fertility clinic. The study population contains men with a range of semen quality, selected to some degree by both fertility potential and access to healthcare, limiting generalizability. Additional studies of




Figure 1 Number of publications per year on '(pesticide) and (semen or sperm)' in PubMed, during 1963–2014 ($N = 1171$).

fertile (i.e. partners of pregnant women) and/or unselected by fertility (i.e. students or military candidates) populations would extend generalizability. Because Chiu and colleagues provided only limited data on patient selection, and on men who did and did not complete the FFQ, it is difficult to address potential selection bias. In addition, these findings need to be replicated in various populations as effects of dietary pesticide exposure on semen quality might be modified by genetic or phenotypic background (Perez-Herrera *et al.*, 2008).

Since 1980 the World Health Organization (WHO) has published a series of laboratory manuals for semen analysis that define the standards in the field, recommending hemocytometer for determination of sperm concentration (WHO, 2010). Chiu *et al.* used computer-aided sperm analysis (CASA) methods to estimate sperm parameters. Until recently, it was not feasible to accurately measure sperm concentration by CASA because of difficulties in distinguishing spermatozoa from particulate debris. However, advances in technology, particularly in the use of fluorescent DNA stains and tail-detection algorithms, may now allow sperm concentration to be estimated by CASA methods, provided that adequate care is taken in preparing specimens and proper quality-control procedures are followed. In this study, ejaculate volume was measured by graduated pipette, a method that has been shown to underestimate the volume and is not recommended (WHO, 2010). This is important since calculation of total sperm count is based on ejaculate volume and sperm concentration.

The authors gained statistical power despite limited sample size ($N = 155$) by using generalized linear mixed models with random intercepts, to account for within-person correlations between repeated samples. Careful selection of covariates (age, body mass index, smoking, abstinence time, physical activity, total energy intake, history of varicocele and dietary pattern) likely also contributed to the precision of their estimates. Thus, despite the relatively small sample size and exposure assessment limitations, the paper make a convincing case that dietary exposure to pesticides can adversely impact semen quality. While this finding will need to be replicated in other settings and populations, it carries important public health implications.

Public health implications

Male fertility, strongly related to poor semen quality, is of considerable public health importance for several reasons. First, its medical, societal and economic burden is high, as it is a leading cause of unsuccessful attempts to achieve pregnancy and one of the most common medical problems among young men (Winters and Walsh, 2014). Second, it has been suggested as an important marker of male health, predicting both morbidity and mortality (Jensen *et al.*, 2009; Eisenberg *et al.*, 2014a,b). Third, it is sensitive to environmental exposures, including endocrine disrupting chemicals, heat and life-style factors, such as diet (Afeiche *et al.*, 2013; Bergman *et al.*, 2013) or body mass index (Eisenberg *et al.*, 2014a,b). Therefore, it can provide a sensitive marker of the impacts of modern environment on human health (Nordkap *et al.*, 2012).

This paper by Chiu and colleagues contributes to the growing body of evidence that non-occupational exposures, and particularly dietary factors (i.e. meat, fish, dairy, etc.) can impact male fertility (Swan *et al.*, 2007; Gaskins *et al.*, 2012; Afeiche *et al.*, 2014a, b). Several prior studies have examined semen quality in relation to dietary pesticide exposure and/or organic diet (Jensen *et al.*, 1996; Juhler *et al.*, 1999). Several studies have shown that substituting an organic for a conventional diet can reduce exposure to pesticides, at least among children (Lu *et al.*, 2006). Currently, the male partner in couples attending fertility clinic rarely receive assessment or recommendation regarding life-style and dietary factors. Lessons learned from well-conducted studies should be considered when advising men attempting to conceive. In addition, there is clear need for further studies on diet and male fertility, including interventional studies. Such studies can be conducted in a range of settings besides the fertility clinic. One particularly promising, though challenging approach, is to enroll the couple at the time of pregnancy planning and assess the exposure and fecundity of both partners (Louis *et al.*, 2015).

Despite its importance, male infertility is not a major focus of the research community (Louis, 2014). This may be due to increasing population growth in some areas leading to the perception that contraception is

the primary public health concern. Moreover, assisted reproductive technology has greatly improved the likelihood of conception despite poor semen quality. However, these techniques are costly, both in resources and health, and there is concern regarding long-term implications of such techniques (Land and Evers, 2003). When fertility is addressed, the male partner is often overlooked; contraception and infertility have been perceived to be primarily female issues, and the root causes of male infertility are not being addressed. These causes might differ along the life course, as male infertility is determined both during fetal development and during adulthood (Juul *et al.*, 2014). As shown by Chiu *et al.*, male fertility studies provide an opportunity not only to better understand the causes of an important public health problem, but more broadly to illustrate effects of the modern environment on human health.

Acknowledgements

Hagai Levine gratefully acknowledges support by a post-doctoral fellowship award from the Environment and Health Fund (EHF), Jerusalem, Israel, and supplementary support from American Healthcare Professionals and Friends for Medicine in Israel (APF) and Israel Medical Association (IMA).

Funding

No external funding was either sought or obtained for this study.

Conflict of interest

The authors declare they have no actual or potential competing financial interests.

References

- Afeiche M, Williams P, Mendiola J, Gaskins A, Jørgensen N, Swan S, Chavarro J. Dairy food intake in relation to semen quality and reproductive hormone levels among physically active young men. *Hum Reprod* 2013;**28**: 2265–2275.
- Afeiche MC, Bridges ND, Williams PL, Gaskins AJ, Tanrikut C, Petrozza JC, Hauser R, Chavarro JE. Dairy intake and semen quality among men attending a fertility clinic. *Fertil Steril* 2014a;**101**:1280–1287.
- Afeiche MC, Gaskins AJ, Williams PL, Toth TL, Wright DL, Tanrikut C, Hauser R, Chavarro JE. Processed meat intake is unfavorably and fish intake favorably associated with semen quality indicators among men attending a fertility clinic. *J Nutr* 2014b;**144**:1091–1098.
- Bergman Å, Heindel JJ, Jobling S, Kidd KA, Zoeller RT, Jobling SK. *State of the Science of Endocrine Disrupting Chemicals 2012: an Assessment of the State of the Science of Endocrine Disruptors Prepared by a Group of Experts for the United Nations Environment Programme and World Health Organization, 2013*. World Health Organization, 2013.
- Berman T, Goldsmith R, Goen T, Spungen J, Novack L, Levine H, Amitai Y, Shohat T, Grotto I. Urinary concentrations of organophosphate pesticide metabolites in adults in Israel: demographic and dietary predictors. *Environ Int* 2013;**60**:183–189.
- Chiu YH, Afeiche MC, Gaskins AJ, Williams PL, Petrozza JC, Tanrikut C, Hauser R, Chavarro JE. Fruit and vegetable intake and their pesticide residues in relation to semen quality among men from a fertility clinic. *Hum Reprod* 2015; doi:10.1093/humrep/dev064.
- Curl CL, Beresford SA, Fenske RA, Fitzpatrick AL, Lu C, Nettleton JA, Kaufman JD. Estimating pesticide exposure from dietary intake and organic food choices: the Multi-Ethnic Study of Atherosclerosis (MESA). *Environ Health Perspect* 2015. doi:10.1289/ehp.1408197. (Epub ahead of print).
- Eisenberg ML, Kim S, Chen Z, Sundaram R, Schisterman EF, Louis GMB. The relationship between male BMI and waist circumference on semen quality: data from the LIFE study. *Hum Reprod* 2014a;**29**:193–200.
- Eisenberg ML, Li S, Behr B, Cullen MR, Galusha D, Lamb DJ, Lipshultz LI. Semen quality, infertility and mortality in the USA. *Hum Reprod* 2014b;**29**:1567–1574.
- Gaskins AJ, Colaci DS, Mendiola J, Swan SH, Chavarro JE. Dietary patterns and semen quality in young men. *Hum Reprod* 2012;**27**:2899–2907.
- Jensen TK, Giwercman A, Carlsen E, Scheike T, Skakkebaek NE. Semen quality among members of organic food associations in Zealand, Denmark. *Lancet* 1996;**347**:1844.
- Jensen TK, Jacobsen R, Christensen K, Nielsen NC, Bostofte E. Good semen quality and life expectancy: a cohort study of 43,277 men. *Am J Epidemiol* 2009;**170**:559–565.
- Juhler RK, Larsen SB, Meyer O, Jensen ND, Spano M, Giwercman A, Bonde JP. Human semen quality in relation to dietary pesticide exposure and organic diet. *Arch Environ Contam Toxicol* 1999;**37**:415–423.
- Juul A, Almstrup K, Andersson A-M, Jensen TK, Jørgensen N, Main KM, Meys ER-D, Toppari J, Skakkebaek NE. Possible fetal determinants of male infertility. *Nat Rev Endocrinol* 2014;**10**:553–562.
- Land J, Evers J. Risks and complications in assisted reproduction techniques: Report of an ESHRE consensus meeting. *Hum Reprod* 2003;**18**:455–457.
- Louis GM. Male fecundity and its implications for health and disease across the lifespan. *Hum Reprod* 2014;**29**:1351–1352.
- Louis GM, Chen Z, Schisterman EF, Kim S, Sweeney AM, Sundaram R, Lynch CD, Gore-Langton RE, Barr DB. Perfluorochemicals and human semen quality: the LIFE study. *Environ Health Perspect* 2015;**123**:57–63.
- Lu C, Toepel K, Irish R, Fenske RA, Barr DB, Bravo R. Organic diets significantly lower children's dietary exposure to organophosphorus pesticides. *Environ Health Perspect* 2006;**114**:260–263.
- Martenies SE, Perry MJ. Environmental and occupational pesticide exposure and human sperm parameters: a systematic review. *Toxicology* 2013;**307**:66–73.
- Nordkap L, Joensen UN, Blomberg Jensen M, Jørgensen N. Regional differences and temporal trends in male reproductive health disorders: semen quality may be a sensitive marker of environmental exposures. *Mol Cell Endocrinol* 2012;**355**:221–230.
- Perez-Herrera N, Polanco-Minaya H, Salazar-Arredondo E, Solis-Heredia MJ, Hernandez-Ochoa I, Rojas-Garcia E, Alvarado-Mejia J, Borja-Aburto VH, Quintanilla-Vega B. PON1Q192R genetic polymorphism modifies organophosphorus pesticide effects on semen quality and DNA integrity in agricultural workers from southern Mexico. *Toxicol Appl Pharmacol* 2008;**230**:261–268.
- Swan SH, Liu F, Overstreet JW, Brazil C, Skakkebaek NE. Semen quality of fertile US males in relation to their mothers' beef consumption during pregnancy. *Hum Reprod* 2007;**22**:1497–1502.
- World Health Organization. *WHO Laboratory Manual for the Examination and Processing of Human Sperm*, 5th edn. Geneva, Switzerland: World Health Organization, 2010.
- Whorton D, Krauss RM, Marshall S, Milby TH. Infertility in male pesticide workers. *Lancet* 1977;**2**:1259–1261.
- Winters BR, Walsh TJ. The epidemiology of male infertility. *Urol Clin North Am* 2014;**41**:195–204.