

Time to take human embryo culture seriously†

Arne Sunde^{1,*}, Daniel Brison², John Dumoulin³, Joyce Harper⁴, Kersti Lundin⁵, M. Cristina Magli⁶, Etienne Van den Abbeel⁷, and Anna Veiga⁸

¹Department of Obstetrics and Gynaecology, St. Olav's University Hospital in Trondheim, Trondheim, Norway ²Department of Reproductive Medicine, St Mary's Hospital, Central Manchester NHS Foundation Trust, Manchester Academic Health Sciences Centre, Manchester, UK ³Department of Obstetrics and Gynaecology, IVF Laboratory, Maastricht University Medical Center, Maastricht, The Netherlands ⁴Embryology, IVF and reproductive genetics group, Institute for Women's Health, University College London, London, UK ⁵Reproductive Medicine, Sahlgrenska University Hospital, Göteborg, Sweden ⁶SISMER, Reproductive Medicine Unit, Bologna, Italy ⁷Reproductive Medicine, Gent University Hospital, Gent, Belgium ⁸Reproductive Medicine Service, Hospital Universitari Dexeus, Barcelona, Spain

*Correspondence address. E-mail: trondheimivf@gmail.com

STUDY QUESTION: Is it important that end-users know the composition of human embryo culture media?

SUMMARY ANSWER: We argue that there is as strong case for full transparency concerning the composition of embryo culture media intended for human use.

WHAT IS KNOWN ALREADY: Published data suggest that the composition of embryo culture media may influence the phenotype of the offspring.

STUDY DESIGN, SIZE, DURATION: A review of the literature was carried out.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Data concerning the potential effects on embryo development of culture media were assessed and recommendations for users made.

MAIN RESULTS AND THE ROLE OF CHANCE: The safety of ART procedures, especially with respect to the health of the offspring, is of major importance. There are reports from the literature indicating a possible effect of culture conditions, including culture media, on embryo and fetal development. Since the introduction of commercially available culture media, there has been a rapid development of different formulations, often not fully documented, disclosed or justified. There is now evidence that the environment the early embryo is exposed to can cause reprogramming of embryonic growth leading to alterations in fetal growth trajectory, birthweight, childhood growth and long-term disease including Type II diabetes and cardiovascular problems. The mechanism for this is likely to be epigenetic changes during the preimplantation period of development. In the present paper the ESHRE working group on culture media summarizes the present knowledge of potential effects on embryo development related to culture media, and makes recommendations.

LIMITATIONS, REASONS FOR CAUTION: There is still a need for large prospective randomized trials to further elucidate the link between the composition of embryo culture media used and the phenotype of the offspring. We do not presently know if the phenotypic changes induced by *in vitro* embryo culture represent a problem for long-term health of the offspring.

WIDER IMPLICATIONS OF THE FINDINGS: Published data indicate that there is a strong case for demanding full transparency concerning the compositions of and the scientific rationale behind the composition of embryo culture media.

STUDY FUNDING/COMPETING INTEREST(S): This work was funded by The European Society for Human Reproduction and Embryology. No competing interests to declare.

Key words: assisted reproductive technology / European Society of Human Reproduction and Embryology / culture media / regulation of culture media / effect of *in vitro* culture on human embryos / transparency regarding composition of culture media

†ESHRE Pages content is not externally peer reviewed. The manuscript has been approved by the Executive Committee of ESHRE.

Introduction

For a successful pregnancy to occur, embryos that are able to implant are essential. Several studies have suggested that the composition of the media that embryos are cultured in may have an impact on the quality of embryos generated in IVF/ICSI cycles thereby influencing implantation and pregnancy rates (Mantikou *et al.*, 2013).

Looking back, the culture of human gametes and embryos has been performed with surprisingly little standardization and a lot of experimentation. In the first decade of human IVF, the culture media ranged from fairly simple salt solutions such as Earle's or Tyrode's solutions to complex media intended for tissue culture such as HamF10. All of them were usually supplemented with different sources of proteins, mostly with fetal or maternal serum. Even today commercially available culture media range from fairly simple with 8 to 10 different salts and sugars to media containing nearly 80 different components including amino acids, lipids, vitamins, trace ions and bioactive molecules such as hormones and expression modulators (Gomes *et al.*, 2009; Sunde and Balaban, 2013; Dyrlund *et al.*, 2014; Morbeck *et al.*, 2014a; Chronopoulou and Harper, 2015).

Today, culture media for embryos can be used either as a sequential system, with different compositions for Days 0–3 and 3–6, or as a single medium used for the whole culture period. Culture media should be used as intended by the manufacturer, however, in many laboratories combinations of media from different manufacturers are used (Gardner and Lane, 1997; Biggers and Summers, 2008), making the evaluation of potential impact more difficult.

It is well documented from epidemiological data that the nutritional status of the mother in early pregnancy may have lasting effects on the health of the offspring (Roseboom *et al.*, 2006; Heijmans *et al.*, 2008; Huang *et al.*, 2010). In animal models, it is demonstrated that minor dietary changes that influence the composition of the intrauterine fluid at the time of implantation will have an effect on the cardiovascular system in the offspring (Kwong *et al.*, 2000; Watkins *et al.*, 2007). These observations have led to the formulation of the Developmental Origins of Health and Disease (DOHaD) hypothesis (Barker, 2007; <https://dohadsoc.org>). This can be understood as a physiological response of the embryo to environmental conditions, most likely mediated through epigenetic changes that may persist throughout life and into subsequent generations.

Following this line of reasoning, it would not be surprising if different culture media have different effects on the epigenome of the developing pre-embryo, and data suggest that *in vitro* culture of gamete and embryos may influence the phenotype of the offspring (Khosla *et al.*, 2001; Gomes *et al.*, 2009; Katari *et al.*, 2009; El Hajj and Haaf, 2013; Nelissen *et al.*, 2014). The consequences of this potential shift in phenotype after IVF is to a large extent unknown, and it is unclear whether this will have an effect on the long-term health of the ART-offspring.

The following sections and Table I summarize the knowledge currently available on the main components of culture media and their possible effects on embryo and perinatal development and long-term outcome.

Media composition and significant changes made by manufacturers

Since 1990s, manufacturers have introduced numerous changes in culture media supplements, with the aim to improve success rates of

IVF/ICSI. As shown in a recent study (Morbeck *et al.*, 2014a), commercial IVF media vary widely in their nutrient composition, so to evaluate possible effects on embryo growth it is crucial to know the composition of the media the embryo has been exposed to. For example, amino acids (e.g. methionine) and vitamins taken up by the embryo have been shown to cause direct epigenetic changes via the one carbon cycle of the embryo (Menezo *et al.*, 2013; Xu and Sinclair, 2015). The quantity of these vitamins and amino acids in culture media is variable and poorly defined.

Another important component is peptide growth factors, which are present in some commercially available IVF media. Growth factors have the ability to alter embryonic growth, cell lineage allocation to inner cell mass (ICM) and trophectoderm (TE), and apoptosis, the mechanism by which developing organisms eliminate cells that contain genomic damage. Embryos cultured in small volumes of medium and/or without changing medium are likely to be exposed to biologically active concentrations of growth factors they secrete themselves (autocrine factors) (O'Neill, 2008).

Another important component in culture media formulations, which is still under debate, is protein/albumin supplementation. Human serum albumin (HSA) and protein preparations contain not only albumin, but also a range of potentially bioactive molecules and contaminants (Morbeck *et al.*, 2014b). We still do not know to what extent different preparations of HSA can affect the development of human embryos *in vitro*. Several questions remain about culture media protein supplementation, such as: Is recombinant albumin as effective as sera-derived albumin or a combination of albumins and globulins? Can hyaluronan replace albumin?

It is still unclear how the composition of the media affects embryo quality and IVF/ICSI success rates and which—if any—culture medium leads to the highest IVF/ICSI success rates. Mantikou *et al.* (2013) performed a systematic review that including 22 studies of 20 different culture media from 11 commercial companies, but was unable to find a superior culture medium from the pooled data. It is important to underline that by considering pregnancy or implantation rates only from fresh cycles, this review did not investigate possible long-term effects of culture media.

The potential impact that varying concentrations of components (such as energy substrates, amino acids, EDTA, macromolecules) and other variables (such as growth factors and osmolality) may have on early embryo development, live birth rates and on long-term health implications in the newborn or adult is not fully known. When comparing the potential impact of culture media formulations on embryo quality as well as on the epigenome, two additional questions must be considered: how long the embryo is cultured *in vitro*, and what culture environment is provided by the laboratory.

For quality control, culture media companies use the mouse embryonic assay (MEA) (Ackerman *et al.*, 1984, 1985; Byers *et al.*, 2006). A specific culture medium is considered to pass the MEA if it supports development of $\geq 80\%$ of mouse embryos to the blastocyst stage. Unfortunately, there is no standardization on the methodology of this assay, i.e. on the day of development the culture should start, the mouse strain to be used, the minimum number of embryos to be cultured, the presence of oil overlay, the supplementation with proteins, osmolality etc. This raises concerns regarding the reliability of the current MEA as a proof of the suitability and safety of culture media for human embryos. No alternative is currently available.

Table I List of the main components present in culture media used in assisted reproductive technology procedures.

	What it is known	References	Conclusions
Carbohydrates	<ul style="list-style-type: none"> - Uptake of pyruvate remains high throughout human preimplantation embryo development. - Shift in preference from pyruvate to glucose around the time of compaction. - Glucose alone does not support embryo development before the eight-cell stage. - Lactate alone does not support embryo development before the two-cell stage. 	<p>Lane and Gardner (2007) Morbeck et al. (2014a,b) Menezo et al. (2013)</p>	<p>Glucose inhibition of embryo development is still a matter of debate. It is now generally accepted that glucose should be included in media at the concentrations found in the human oviduct.</p>
Amino acids	<ul style="list-style-type: none"> - Precursors of biosynthesis, energy source, regulator, buffer, chelator. - Non-essential amino acids (alanine, asparagine, aspartate, glutamate, glycine, proline and serine) and glutamine stimulate cleavage rate, blastocyst formation and hatching. - Essential amino acids stimulate cleavage rates after the eight-cell stage and increase development of the inner cell mass in the blastocyst. - Oviduct fluid contains all 20 amino acids. - Follicular fluid and oviduct fluid have concentrations of amino acids much lower than that present in sequential media formulations. 	<p>Lane and Gardner (2007) Morbeck et al. (2014a,b) Menezo et al. (2013)</p>	<ul style="list-style-type: none"> - The importance of amino acids is attributed to their roles in protein and nucleotide synthesis, in acting as osmolytes, as precursors for signaling molecules, and as substrates for energy metabolism. - No currently used media contain optimal concentrations of amino acids. - Glutamine can be replaced with stable glycyl-glutamine to avoid breakdown to products including ammonium.
EDTA	<ul style="list-style-type: none"> - Sequential media formulations usually utilize a concentration of EDTA around 0.01–0.1 mmol/l. - For single step media, the presence of EDTA for the entire culture period has no significant affect on pre- or post-implantation development provided it is used at a low concentration. 	<p>Lane and Gardner (2007) Morbeck et al. (2014a,b) Menezo et al. (2013)</p>	<ul style="list-style-type: none"> - EDTA supplementation is still a matter of debate: it has been suggested that EDTA concentrations should be high in order to regulate glycolysis, but other studies using lower concentrations of EDTA in culture media do not support this statement.
Protein supplementation	<ul style="list-style-type: none"> - Albumin can be a carrier for lipids, hormones, and minerals and can also be associated with globulins. - The two most common commercial protein supplements in human IVF are human serum albumin and human serum albumin with globulins. - There are still concerns about commercially prepared blood-derived albumin and albumin/globulin. 	<p>Lane and Gardner (2007) Morbeck et al. (2014a,b) Menezo et al. (2013) Chronopoulou and Harper (2015)</p>	<ul style="list-style-type: none"> - Human serum albumin (HSA) preparations contain not only albumin, but also a range of potentially bioactive molecules. - It is still unknown to what extent heterogeneity in different preparations of HSA affects the development of human embryos <i>in vitro</i>. - Hyaluran has been suggested as being advantageous for transfer and embryo implantation, but the reasoning for this is still unclear.
Osmolality	<ul style="list-style-type: none"> - The physiological osmolarity surrounding the embryo <i>in vivo</i> is 280–290 mOsm/kg. - Most of the culture media today are formulated to fall in a range of 275–290 mOsm. - There is no evidence of altered development when using media of lower osmolality (= 260 mOsm). 	<p>Brinster (1965) Menezo et al. (2013) Biggers and Racowsky (2001)</p>	<ul style="list-style-type: none"> - The optimal osmolality for human embryo culture is still unknown.

Continued

Table 1 Continued

What it is known	References	Conclusions
<p>Growth factors and cytokines</p> <ul style="list-style-type: none"> - Growth factors have the ability to alter embryonic growth, cell lineage allocation to inner cell mass and trophectoderm, and apoptosis. - A variety of growth factors and cytokines are present in the female reproductive tract. - Most of them are also produced and secreted by the embryo, but not granulocyte-macrophage colony-stimulating factor (GM-CSF). - Insulin added in high doses by some media manufacturers. - GM-CSF added to media by one manufacturer. 	<p>Lane and Gardner (2007) Morbeck <i>et al.</i> (2014a,b), Menezo <i>et al.</i> (2013) Chronopoulou and Harper (2015) Morbeck <i>et al.</i> (2014a,b)</p>	<ul style="list-style-type: none"> - Embryos are likely to benefit from a cocktail of growth factors at optimum concentration. - However, as growth factors act in both positive and negative synergy in order to produce an effect, addition of a single growth factor is questionable. - Growth factors, if not well regulated, could have adverse effects on embryo development and have been associated with large offspring syndrome. - The safe addition to culture media of single or mixtures of growth factors has not been reached yet. - Embryos themselves secrete autocrine factors, whose effect is especially marked when cultured in small volumes of medium and/or without changing medium.
<p>Vitamins, nucleotides, steroids</p> <ul style="list-style-type: none"> - Vitamins are taken up by the embryo and have the potential to cause direct epigenetic changes via the I carbon cycle. - Some media contain vitamins in variable and poorly defined quantities. 	<p>Menezo <i>et al.</i> (2013) Xu and Sinclair (2015)</p>	<ul style="list-style-type: none"> - The role of vitamins, nucleotides and steroids during early development is unclear.

Impact of ART culture medium on embryonic development

The genome and epigenome of the embryo are completely remodeled (integration of the paternal genome, transition from maternal to zygotic transcripts, global genome demethylation and remethylation, maintenance of genetic imprints marking parent of origin alleles) between fertilization and implantation, making the embryo uniquely vulnerable at precisely the stage of manipulation *in vitro* and culture during IVF. At the blastocyst stage, changes in the allocation of cells to the ICM and TE lineages also occur. This can alter ICM/fetal size directly or indirectly via supply of nutrients by altering TE and hence placental size and function, and thereby altering fetal growth trajectory and birthweight.

Culture media contain a number of components, which might affect epigenetic or other reprogramming events, leading to detrimental effects on the embryo and the newborn, and potentially to long-term health problems in the adult. A main component is nutrients. There is evidence that cells—including early embryos—undergo ‘nutrient sensing’ in order to determine the environment in which they are developing. Glucose and amino acids are particularly important, triggering responses via the 5’ adenosine monophosphate-activated protein kinase and mechanistic target of rapamycin (mTOR) pathways, both of which regulate growth and development including via epigenetic changes. Embryonic sensing of a low nutrient environment in IVF culture media can therefore program the embryo to be ‘thrifty’ in its metabolism, programming a long-term thrifty phenotype. When the embryo is then replaced in the maternal environment, it encounters a high nutrition environment and experiences overgrowth (Eckert *et al.*, 2015; Pantaleon, 2015). The reverse would also be true in a high nutrient culture medium environment.

The background literature in animals is extensive, showing that nutrition during preimplantation development, either via the maternal tract or in culture, can cause long-term reprogramming of development and health (Kwong *et al.*, 2000; Eckert *et al.*, 2015). This is likely mediated by epigenetic changes, and there is good evidence that embryo culture particularly in suboptimal conditions disrupts the maintenance of the embryonic epigenome in the mouse and potentially in human preimplantation embryos (Market-Velker *et al.*, 2010; Velker *et al.*, 2012; White *et al.*, 2015).

The impact of culture medium can be captured by global gene expression profiling (transcriptomics) of individual embryos (e.g. mouse: Rinaudo and Schultz, 2004; bovine: Urrego *et al.*, 2014). Human embryos show intrinsic heterogeneity in their transcriptomes (Vassena *et al.*, 2011; Shaw *et al.*, 2013) with evidence of an impact of culture medium (Kleijkers *et al.*, 2015a). Cryopreserved human embryos also show a different gene expression pattern to fresh embryos, including genes regulating formation of ICM and TE, suggesting a possible mechanism for altered fetal growth and placental function in children arising from cryopreserved embryos (Shaw *et al.*, 2012).

Potential impact of culture media on children

It has been shown in several systematic reviews and meta-analyses that singletons born after IVF or ICSI are at increased risk of adverse perinatal

outcome when compared with naturally conceived children with respect to preterm birth, lower birthweight, perinatal mortality and congenital abnormalities (Helmerhorst et al., 2004; Jackson et al., 2004; McDonald et al., 2009; Pandey et al., 2012; Wen et al., 2012; Feuer et al., 2013; Pinborg et al., 2013; Albertini et al., 2014; Lazaraviciute et al., 2014; Marino et al., 2014; Simpson, 2014; Declercq et al., 2015).

According to the DoHAD hypothesis, birthweight is an indication of altered fetal growth and programming of later life health and disease via intermediate physiology. Thus, it is significant that in addition to lower ART birthweights, a growing number of studies have shown distinct phenotypic differences, including those predicted by DoHAD, between ART children and naturally conceived children from birth until young adult age with respect to anthropometric parameters and adiposity (Green et al., 2013), cardiometabolic characteristics such as blood pressure and cardiovascular remodeling (Ceelen et al., 2009; Scherrer et al., 2012; Valenzuela-Alcaraz et al., 2013; Yeung and Druschel, 2013; Chen et al., 2014; Zhou et al., 2014; Pontessilli et al., 2015) and metabolites linked to obesity, insulin resistance and metabolic syndrome (Green et al., 2013; Gkourogianni et al., 2014; Lou et al., 2014). However, if there is evidence regarding an altered birthweight in ART children and some phenotypic differences, several years have to pass until the children age, to establish possible associations with the actual disease.

Although this growing list of studies reporting adverse perinatal and long-term outcome parameters of ART children shows compelling evidence that there could be an association between the mode of conception and outcome, there are several important points to consider. First, it must be recognized that this association may be affected by bias and confounding factors, in particular in the selection of control groups. ART offspring need careful surveillance and follow up in order to record and evaluate possible adverse effects of the treatments (Barnhart, 2013). Second, it must be realized that some reported phenotypic differences are in the physiological range and, most importantly, that the implications are unknown. There are a number of publications that demonstrate that long-term health of ART children is comparable to that of spontaneously conceived children (see the extensive reviews of Bay et al., 2013, 2014; Hart and Norman, 2013a,b; Hediger et al., 2013; Fauser et al., 2014; Halliday et al., 2014). Third, even if the assumption of an association of ART and adverse outcomes in the children is true, there is no evidence that the origin of the problem is the ART treatment itself. Patient-related factors such as the subfertility of the progenitors and associated genetic factors may be one reason for adverse perinatal and post-natal outcomes in singletons after ART. There are reports showing that patient-related factors may be responsible for the differences observed in birthweight and other perinatal outcomes in ART children (Romundstad et al., 2008; Raatikainen et al., 2012; Messerlian et al., 2013; Declercq et al., 2015; Wise et al., 2015). For certain outcome parameters, such as the rate of congenital abnormalities, patient-related factors are considered to be the most important (Simpson, 2014).

However, several studies show that specific ART treatment related factors are at least in part responsible for the observed differences in pregnancy outcome (see extensive review by Pinborg et al., 2013). For instance, in several studies that compare children born after fresh transfer with those born after transfer of frozen/thawed embryos, it was shown that singletons born after frozen embryo transfer have a significantly higher mean birthweight and higher risk of large for gestational age, macrosomia and perinatal mortality when compared with singletons conceived after fresh transfer (Henningsen et al., 2011; Nakashima

et al., 2013; Wennerholm et al., 2013; Marino et al., 2014). Here, subfertility could be ruled out as a determining factor for these differences, since children from both groups were born from the same population of subfertile women. It is not clear from these studies which aspects of the ART technology (hormonal stimulation or cryopreservation) are responsible for the observed differences. Interestingly it was found that human embryos surviving the cryopreservation and thawing procedure exhibit a different gene expression compared with fresh embryos matched for morphological grade (Shaw et al., 2012).

ART children have also shown alterations in DNA methylation and transcript level of a number of genes such as those controlling growth (insulin-like growth factor (IGF)2/H19 and its receptor, IGF2R) in cord blood and placenta (Katari et al., 2009; Turan et al., 2010; Zhang et al., 2010; Song et al., 2015). The most recent of these studies strongly suggests that the causal effect is from ART procedures, rather than from parental infertility and may provide insight into the reported increases in imprinting disorders in ART children (Song et al., 2015).

Whether or not the *in vitro* culture of embryos, and more specifically, the culture medium used (or certain components of media), is also a determining factor for the observed adverse pregnancy outcome after ART was recently discussed in a review by Zandstra et al. (2015). It was concluded that from the 11 studies that investigated the association between birthweight after human IVF and the type of culture medium used, 5 found evidence of such an association while the other 6 studies did not. From these findings, together with the fact that the number of human studies is still limited, and that most studies were retrospective with consecutive use of different culture media and limited sample sizes, making bias of results likely, it can be concluded that it is still unclear whether or not culture media (or components in these media) can affect outcome parameters such as birthweight. Furthermore, it is unknown whether or not any possible relationship with birthweight is restricted to certain media or specific components. Interestingly, it was demonstrated in a recent study that birthweight of singletons born after IVF showed an inverse association with the storage time of the embryo culture medium used, indicating a possible accumulation of deterioration of certain medium components (Kleijkers et al., 2015b).

If a relationship between the culture medium used and certain perinatal outcome parameters such as birthweight is found to be true, it is still too early to know the long-term effect on health. In one recent study it was found that the differences in birthweight as a result of the type of culture medium used persisted during the first 2 years of life (Kleijkers et al., 2014).

Regulatory aspects of ART culture media

In 2009 it was agreed that the European Union (EU) would regulate ART culture media as medical devices (Class III) (Medical Device Directive 93/42/EEC, 1993; MEDDEV2.2/4 Jan, 2012).

CE (Conformité Européenne) marking is a regulatory requirement to show a product is safe, effective, fit for purpose and meets EU regulations. To comply with European Directive 2001/83/EC, culture medium should be preferably CE marked ('Wherever possible, only CE marked medical devices must be used and all concerned staff must have received appropriate training on the use of such devices' Commission Directive, 2006/17). To apply for CE marking, the manufacturers

have to submit a technical file to their Notified Body (an organization that has been accredited by a Member State to assess whether a product meets certain preordained standards) which should include: the grounds for incorporating each component in the media, the risk of each of these components, manufacturing processes, design, validation of design, testing of product, toxicology, scientific data, what international standards have been used, pre-clinical and post clinical evaluation, ensure the product will not compromise safety of patients, and errors and a risk benefit analysis. The clinical evaluation must show that there is an overall positive risk/benefit ratio of the product and take into account the safety to the gametes, embryos and the mother to be (in accordance with [MEDDEV 2.7-1](#)). The Notified Body will also do a physical audit of the production site and look at the quality management system and manufacturing process, post market surveillance and follow up. Manufacturers should also have a plan for post-market surveillance including clinical follow-up (in accordance with [MEDDEV 2.12/1](#)).

If the medium contains a medicinal product or human derived blood product, such as HSA or antibiotics, the Notified Body also has to undertake a consultation with a Medicines Competent Authority and/or European Medicines Agency. In both cases, they will look at safety, performance and usefulness of the element within the media.

If the manufacturer makes any changes to a medium, they would need to have these changes validated and added to their CE certificate after review by their Notified Body and completion of all relevant consultations ([Medical Device Directive 93/42/EEC, 1993](#)).

Currently, it is not clear how the Notified Body system works. As far as we are aware, it is not possible to find out who the notified bodies are, their membership, details of advisors or relevant expertise. It is also currently unclear how manufacturers carry out post-market surveillance and clinical follow-up arising from use of culture medium in IVF.

As stated above, culture media should be used as intended by the manufacturer, as an incorrect use can invalidate the CE marking.

Recommendations for end users

For IVF laboratories, the use of devices and equipment, including culture media, is a constant decision-making process, which has to be made on the basis of a number of criteria such as availability, user friendliness and clinical results, as well as cost/benefit. The clinics have responsibility for the products that they use and decision-making processes have to be based on available information regarding the characteristics of the product itself and its suitability, including possible changes. In addition, since culture media manufacturers often make changes (minor or major) in their products, decisions have to be re-evaluated and re-validated.

End users must be aware of when changes are being made. It is therefore extremely important that culture media manufacturers notify clinics of all changes being made to their products. Changes in the formulation and other characteristics of the culture medium have to be declared, and the scientific basis of the change has to be disclosed, including validation in experimental models (animal models) as well as clinical results, if available. The expected consequences of the medium change have to be discussed, especially with respect to safety concerns.

In parallel, the clinics should have a system to validate the products they use in their own laboratory. The results obtained with new media or other methodological changes must be carefully monitored and

assessed. Multicentre studies that will allow a faster validation process of a specific product change should be encouraged.

A quality management system must be in place to ensure the correct methodology of storage and the use of a specific culture medium by all personnel involved according to the manufacturer's instructions. It is mandatory that traceability, including type of culture media and lot number, is maintained for all ART products. In case incidents occur, they must be reported as notification of serious adverse reactions and events ([Commission Directive, 2006/86/EC](#)). Biovigilance systems put in place by the competent authorities should assist in the detection of possible adverse events or reactions that may arise.

Results regarding fertilization rate and embryo quality as well as pregnancy, implantation and live born baby rates have to be carefully recorded and analyzed. Each IVF unit should develop a system for following up also the health of the offspring and this should be one of the crucial quality control indicators. The follow-up of children needs to be performed periodically, preferably through national registries.

In light of the large and emerging body of evidence on the importance of the nutritional environment during early development, national registries should include the type of culture medium embryos have been exposed to as well as unique identifiers for patients and babies on the register to enable large-scale outcome studies to be performed routinely using data linkage.

Conclusions

The ultimate goal of any ART treatment must be to secure the health of the patients and the children. This can only be fulfilled by continuous follow-up and by documentation and traceability of all methods performed and all materials used. As shown in this summary, culture media have a high degree of variability, and in general their quantitative composition is not known by the end users, which limits the possibility to correlate culture media type and composition to outcome.

Existing data regarding the influence of culture media on embryonic, obstetrical, perinatal and post-natal parameters are conflicting. However, the fact that some studies do find such an association, particularly with birthweight, highlights the importance of further investigation.

Therefore, clinical data related to culture media should be documented by ART centers in such a way that they can be used for biovigilance purposes and these data should preferably be collected in national registries, where available.

In addition, the complete formulation of culture media should be disclosed and all changes should be justified, validated and communicated to end-users by manufacturers. Changes should be made to the existing regulatory system to achieve transparency and improve monitoring of outcomes to the long-term benefit of ART children and society.

Authors' roles

All authors participated equally in conceiving, writing and editing the manuscript.

Funding

This work was funded by the European Society for Human Reproduction and Embryology.

Conflict of interest

None declared.

References

- Ackerman SB, Stokes GL, Swanson RJ, Taylor SP, Fenwick L. Toxicity testing for human *in vitro* fertilization programs. *J In Vitro Fert Embryo Transfer* 1985; **2**:132–137.
- Ackerman SB, Taylor SP, Swanson RJ, Laurell LH. Mouse embryo culture for screening in human IVF. *Arch Androl* 1984; **12**:129–136.
- Albertini DF, Evers JLH, Geraedts JPM, Gianaroli L, Sharpe R, Sinclair KD, Sunde A, Van Steirteghem A. (ESHRE Capri Workshop Group) Birth defects and congenital health risks in children conceived through assisted reproduction technology (ART): a meeting report. *J Assist Reprod Genet* 2014; **31**:947–958.
- Barker DJ. The origins of the developmental origins theory. *J Intern Med* 2007; **261**:412–417.
- Barnhart KT. Assisted reproductive technologies and perinatal morbidity: interrogating the association. *Fertil Steril* 2013; **99**:299–302.
- Bay B, Mortensen EL, Kesmodel US. Assisted reproduction and child neurodevelopmental outcomes: a systematic review. *Fertil Steril* 2013; **100**:844–853.
- Bay B, Mortensen EL, Kesmodel US. Is subfertility or fertility treatment associated with long-term growth in the offspring? A cohort study. *Fertil Steril* 2014; **102**:1117–1123.
- Biggers J, Racowsky C. The development of fertilized human ova to the blastocyst stage in medium KSOM-AA: is a two-step protocol necessary? *Reprod Biomed Online* 2001; **5**:133–140.
- Biggers JD, Summers MC. Choosing a culture medium: making informed choices. *Fertil Steril* 2008; **90**:473–483.
- Brinster RL. Studies on the development of mouse embryos *in vitro*. I. The effect of osmolarity and hydrogen ion concentration. *J Exp Zool* 1965; **158**:49–57.
- Byers S, Payson S, Taft R. Performance of ten inbred mouse strains following assisted reproductive technologies (ARTs). *Theriogenology* 2006; **65**:1716–1726.
- Ceelen M, van Weissenbruch MM, Prein J, Smit JJ, Vermeiden JP, Spreuuenberg M, van Leeuwen FE, Delemarre-van de Waal HA. Growth during infancy and early childhood in relation to blood pressure and body fat measures at age 8–18 years of IVF children and spontaneously conceived controls born to subfertile parents. *Hum Reprod* 2009; **24**:2788–2795.
- Chen M, Wu L, Zhao J, Wu F, Davies MJ, Wittert GA, Norman RJ, Robker RL, Heilbronn LK. Altered glucose metabolism in mouse and humans conceived by IVF. *Diabetes* 2014; **63**:3189–3198.
- Chronopoulou E, Harper JC. IVF culture media: past, present and future. *Hum Reprod Update* 2015; **21**:39–55.
- Commission Directive 2006/17/EC of 8 February 2006 implementing Directive 2004/23/EC of the European Parliament and of the Council as regards certain technical requirements for the donation, procurement and testing of human tissues and cells.
- Commission Directive 2006/86/EC of 24 October 2006 implementing Directive 2004/23/EC of the European Parliament and of the Council as regards traceability requirements, notification of serious adverse reactions and events and certain technical requirements for the coding, processing, preservation, storage and distribution of human tissues and cells.
- Declercq E, Luke B, Belanoff C, Cabral H, Diop H, Gopal D, Hoang L, Kotelchuck M, Stern JE, Hornstein MD. Perinatal outcomes associated with assisted reproductive technology: the Massachusetts Outcomes Study of Assisted Reproductive Technologies (MOSART). *Fertil Steril* 2015; **103**:888–895.
- Dyrlund TF, Kirkegaard K, Poulsen ET, Sanggaard KW, Hindkjaer JJ, Kjems J, Enghild JJ, Ingerslev HJ. Unconditioned commercial embryo culture media contain a large variety of non-declared proteins: a comprehensive proteomics analysis. *Hum Reprod* 2014; **29**:2421–2430.
- Eckert JJ, Velazquez MA, Fleming TP. Cell signalling during blastocyst morphogenesis. *Adv Exp Med Biol* 2015; **843**:1–21.
- El Hajj N, Haaf T. Epigenetic disturbances in *in vitro* cultured gametes and embryos: implications for human assisted reproduction. *Fertil Steril* 2013; **99**:632–641.
- Fauser BCJM, Devroey P, Diedrich K, Balaban B, Bonduelle M, Delemarre-van de Waal HA, Estella C, Ezcurra D, Geraedts JPM, Howles CM et al. Health outcomes of children born after IVF/ICSI: a review of current expert opinion and literature. *Reprod BioMed Online* 2014; **28**:162–182.
- Feuer SK, Camarano L, Rinaudo PF. ART and health: clinical outcomes and insights on molecular mechanisms from rodent studies. *Molec Hum Reprod* 2013; **19**:189–204.
- Gardner DK, Lane M. Culture and selection of viable blastocysts: a feasible proposition for human IVF? *Hum Reprod Update* 1997; **3**:367–382.
- Gkourogianni A, Kosteria I, Telonis AG, Margeli A, Mantzou E, Konsta M, Loutradis D, Mastorakos G, Papassotiropou I, Klapa MI et al. Plasma metabolomic profiling suggests early indications for predisposition to latent insulin resistance in children conceived by ICSI. *PLoS One* 2014; **9**:e94001.
- Gomes MV, Huber J, Ferriani RA, Amaral Neto AM, Ramos ES. Abnormal methylation at the KvDMR1 imprinting control region in clinically normal children conceived by assisted reproductive technologies. *Molec Hum Reprod* 2009; **15**:471–477.
- Green MP, Mouat F, Miles HL, Hopkins SA, Derraik JGB, Hofman PL, Peek JC, Cutfield WS. Phenotypic differences in children conceived from fresh and thawed embryos in *in vitro* fertilization compared with naturally conceived children. *Fertil Steril* 2013; **99**:1898–1904.
- Halliday J, Wilson C, Hammarberg K, Doyle LW, Bruinsma F, McLachlan R, McBain J, Berg T, Fisher JR, Amor D. Comparing indicators of health and development of singleton young adults conceived with and without assisted reproductive technology. *Fertil Steril* 2014; **101**:1055–1063.
- Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment. Part I—General health outcomes. *Hum Reprod Update* 2013a; **19**:244–250.
- Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment. Part II—Mental health and development outcomes. *Hum Reprod Update* 2013b; **19**:232–243.
- Hediger ML, Bell EM, Druschel CM, Buck Louis GM. Assisted reproductive technologies and children's neurodevelopmental outcomes. *Fertil Steril* 2013; **99**:311–317.
- Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES, Slagboom PE, Lumey LH. Persistent epigenetic differences associated with prenatal exposure to famine in humans. *Proc Natl Acad Sci USA* 2008; **105**:17046–17049.
- Helmerhorst FM, Perquin DA, Donker D, Keirse MJ. Perinatal outcome of singletons and twins after assisted conception: a systematic review of controlled studies. *BMJ* 2004; **328**:261.
- Henningsen A-K, Pinborg A, Lidegaard O, Vestergaard C, Lyng Forman J, Nyboe Andersen A. Perinatal outcome of singleton siblings born after assisted reproductive technology and spontaneous conception: Danish national sibling-cohort study. *Fertil Steril* 2011; **95**:959–963.
- Huang C, Li Z, Wang M, Martorell R. Early life exposure to the 1959–1961 Chinese famine has long-term health consequences. *J Nutr* 2010; **140**:1874–1878.

- Jackson RA, Gibson KA, Wu YW, Croughan MS. Perinatal outcomes in singletons following *in vitro* fertilization: a meta-analysis. *Obstet Gynecol* 2004;**103**:551–563.
- Katari S, Turan N, Bibikova M, Erinle O, Chalian R, Foster M, Gaughan JP, Coutifaris C, Sapienza C. DNA methylation and gene expression differences in children conceived *in vitro* or *in vivo*. *Hum Molec Genet* 2009;**18**:3769–3778.
- Khosla S, Dean W, Reik W, Feil R. Epigenetic and experimental modification in early mammalian development: Part II. Culture of preimplantation embryos and its long-term effects on gene expression and phenotype. *Hum Reprod Update* 2001;**7**:419–427.
- Kleijkers SH, van Montfoort AP, Smits LJ, Viechtbauer W, Roseboom TJ, Nelissen EC, Coonen E, Derhaag JG, Bastings L, Schreurs IE et al. IVF culture medium affects post-natal weight in humans during the first 2 years of life. *Hum Reprod* 2014;**29**:661–669.
- Kleijkers SH, Eijssen LM, Coonen E, Derhaag JG, Mantikou E, Jonker MJ, Mastenbroek S, Repping S, Evers JL, Dumoulin JC et al. Differences in gene expression profiles between human preimplantation embryos cultured in two different IVF culture media. *Hum Reprod* 2015a;**30**:2303–2311.
- Kleijkers SH, van Montfoort AP, Smits LJ, Coonen E, Derhaag JG, Evers JL, Dumoulin JC. Age of G-1 PLUS v5 embryo culture medium is inversely associated with birthweight of the newborn. *Hum Reprod* 2015b;**30**:1352–1357.
- Kwong WY, Wild AE, Roberts P, Willis AC, Fleming TP. Maternal undernutrition during the preimplantation period of rat development causes blastocyst abnormalities and programming of postnatal hypertension. *Development* 2000;**127**:4195–4202.
- Lane M, Gardner D. Embryo culture medium: which is the best? *Best Pract Res Clin Obstet Gynaecol* 2007;**21**:83–100.
- Lazaravičiute G, Kausar M, Bhattacharya S, Haggarty P, Bhattacharya S. A systematic review and meta-analysis of DNA methylation levels and imprinting disorders in children conceived by IVF/ICSI compared with children conceived spontaneously. *Hum Reprod Update* 2014;**20**:840–852.
- Lou H, Le F, Zheng Y, Li L, Wang L, Wang N, Zhu Y, Huang H, Jin F. Assisted reproductive technologies impair the expression and methylation of insulin-induced gene I and sterol regulatory element-binding factor I in the fetus and placenta. *Fertil Steril* 2014;**101**:974–980.
- Mantikou E, Youssef M, van Wely M, Van der Veen F, Al-inany H, Repping S, Mastenbroek S. Embryo culture media and IVF/ICSI success rates: a systematic review. *Hum Reprod Update* 2013;**19**:210–220.
- Marino JL, Moore VM, Willson KJ, Rumbold A, Whitrow MJ, Giles LC, Davies MJ. Perinatal outcomes by mode of assisted conception and sub-fertility in an Australian data linkage cohort. *PLoS One* 2014;**9**:e80398.
- Market-Velker BA, Fernandes AD, Mann MR. Side-by-side comparison of five commercial media systems in a mouse model: suboptimal *in vitro* culture interferes with imprint maintenance. *Biol Reprod* 2010;**83**:938–950.
- McDonald SD, Han Z, Mulla S, Murphy KE, Beyene J, Ohlsson A. Preterm birth and low birth weight among *in vitro* fertilization singletons: a systematic review and meta-analyses. *Eur J Obstet Gynecol Reprod Biol* 2009;**146**:138–148.
- MEDDEV 93/42/EEC (1993) EU Council Medical Device Directive: <http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=CONSLEG:1993L0042:20071011:en:PDF> (2016, date last accessed).
- MEDDEV 2.2/4 Guidelines for conformity assessment of In Vitro Fertilisation (IVF) and Assisted Reproduction Technologies (ART) products (guidelines that relate to CE marking of culture media): http://ec.europa.eu/health/medical-devices/files/meddev/2_2_4_en.pdf
- MEDDEV 2.7.1(2003) Evaluation of clinical data: A guide for manufacturers and notified bodies
- MEDDEV 2.12-1 rev 8 Guidelines on a Medical Devices Vigilance System (guidance on vigilance with specific reference to ART): http://ec.europa.eu/health/medical-devices/files/meddev/2_12_1_en.pdf (2016, date last accessed).
- Menezes Y, Lichtblau I, Elder K. New insights into human pre-implantation metabolism *in vivo* and *in vitro*. *J Assist Reprod Genet* 2013;**30**:293–303.
- Messerlian C, Maclagan L, Basso O. Infertility and the risk of adverse pregnancy outcomes: a systematic review and meta-analysis. *Hum Reprod* 2013;**28**:125–137.
- Morbeck DA, Krisher R, Herrick J, Baumann N, Matern D, Moyer T. Composition of commercial media used for human embryo culture. *Fertil Steril* 2014a;**102**:759–766.
- Morbeck DE, Paczkowski M, Fredrickson JR, Krisher RL, Hoff HS, Baumann NA, Moyer T, Matern D. Composition of protein supplements used for human embryo culture. *J Assist Reprod Genet* 2014b;**31**:1703–1711.
- Nakashima A, Araki R, Tani H, Ishihara O, Kuwahara A, Irahara M, Yoshimura Y, Kuramoto T, Saito H, Nakaza A et al. Implications of assisted reproductive technologies on term singleton birth weight: an analysis of 25,777 children in the national assisted reproduction registry of Japan. *Fertil Steril* 2013;**99**:450–455.
- Nelissen EC, Dumoulin JC, Busato F, Ponger L, Eijssen LM, Evers JL, Tost J, van Montfoort AP. Altered gene expression in human placentas after IVF/ICSI. *Hum Reprod* 2014;**29**:2821–2831.
- O'Neill C. The potential roles for embryotrophic ligands in preimplantation embryo development. *Hum Reprod Update* 2008;**14**:275–288.
- Pandey S, Shetty A, Hamilton M, Bhattacharya S, Maheshwari A. Obstetric and perinatal outcomes in singleton pregnancies resulting from IVF/ICSI: a systematic review and meta-analysis. *Hum Reprod Update* 2012;**18**:485–503.
- Pantaleon M. The role of hexosamine biosynthesis and signaling in early development. *Adv Exp Med Biol* 2015;**843**:53–76.
- Pinborg A, Aaris Henningsen A-K, Malchau SS, Loft A. Congenital anomalies after assisted reproductive technology. *Fertil Steril* 2013;**99**:327–332.
- Pontessilli M, Painter RC, Grooten IJ, Van der Post JA, Mol BW, Vrijkotte TGM, Repping S, Roseboom TJ. Subfertility and assisted reproduction techniques are associated with poorer cardiometabolic profiles in childhood. *Reprod Biomed Online* 2015;**30**:258–267.
- Raatikainen R, Kuivasaari-Pirinen P, Hippelainen M, Heinonen S. Comparison of the pregnancy outcomes of subfertile women after infertility treatment and in naturally conceived pregnancies. *Hum Reprod* 2012;**27**:1162–1169.
- Rinaudo P, Schultz RM. Effects of embryo culture on global pattern of gene expression in preimplantation mouse embryos. *Reproduction* 2004;**128**:301–311.
- Romundstad LB, Romundstad PR, Sunde A, von Düring V, Skjærven R, Gunnell D, Vatten LJ. Effects of technology or maternal factors on perinatal outcome after assisted fertilisation: a population-based cohort study. *Lancet* 2008;**372**:737–743.
- Roseboom T, de Rooij S, Painter R. The Dutch famine and its long-term consequences for adult health. *Early Hum Dev* 2006;**82**:485–491.
- Scherrer U, Rimoldi SF, Rexhaj E, Stuber T, Duplain H, Garcin S, de Marchi SF, Nicod P, Germond M, Allemann Y et al. Systemic and pulmonary vascular dysfunction in children conceived by assisted reproductive technologies. *Circulation* 2012;**125**:1890–1896.
- Shaw L, Sneddon SF, Brison DR, Kimber SJ. Comparison of gene expression in fresh and frozen-thawed human preimplantation embryos. *Reproduction* 2012;**144**:569–582.
- Shaw L, Sneddon SF, Zeef L, Kimber SJ, Brison DR. Global gene expression profiling of individual human oocytes and embryos demonstrates heterogeneity in early development. *PLoS One* 2013;**22**:e64192.
- Simpson JL. Birth defects and assisted reproductive technologies. *Semin Fetal Neonatal Med* 2014;**19**:177–182.

- Song S, Ghosh J, Mainigi M, Turan N, Weinerman R, Truongcao M, Coutifaris C, Sapienza C. DNA methylation differences between *in vitro*- and *in vivo*-conceived children are associated with ART procedures rather than infertility. *Clin Epigenet* 2015;**7**:41.
- Sunde A, Balaban B. The assisted reproductive technology laboratory: toward evidence-based practice? *Fertil Steril* 2013;**100**:310–318.
- Turan N, Katari S, Gerson LF, Chalian R, Foster MW, Gaughan JP, Coutifaris C, Sapienza C. Inter- and intra-individual variation in allele-specific DNA methylation and gene expression in children conceived using assisted reproductive technology. *PLoS Genet* 2010;**6**:e1001033.
- Urrego R, Rodriguez-Osorio N, Niemann H. Epigenetic disorders and altered gene expression after use of assisted reproductive technologies in domestic cattle. *Epigenetics* 2014;**9**:803–815.
- Valenzuela-Alcaraz B, Crispi F, Bijmens B, Cruz-Lemini M, Creus M, Sitges M, Bartrons J, Civico S, Balasch J, Gratacós E. Assisted reproductive technologies are associated with cardiovascular remodeling *in utero* that persists postnatally. *Circulation* 2013;**128**:1442–1450.
- Vassena R, Boué S, González-Roca E, Aran B, Auer H, Veiga A, Izpisua Belmonte JC. Waves of early transcriptional activation and pluripotency program initiation during human preimplantation development. *Development*. 2011;**138**:3699–3709.
- Velker BA, Denomme MM, Mann MR. Embryo culture and epigenetics. *Methods Mol Biol* 2012;**912**:399–421.
- Watkins AJ, Platt D, Papenbrock T, Wilkins A, Eckert JJ, Kwong WY, Osmond C, Hanson M, Fleming TP. Mouse embryo culture induces changes in postnatal phenotype including raised systolic blood pressure. *Proc Natl Acad Sci USA* 2007;**104**:5449–5454.
- Wen J, Jiang J, Ding C, Dai J, Liu Y, Xia Y, Liu J, Hu Z. Birth defects in children conceived by *in vitro* fertilization and intracytoplasmic sperm injection: a meta-analysis. *Fertil Steril* 2012;**97**:1331–1337.
- Wennerholm UB, Henningsen AK, Romundstad LB, Bergh C, Pinborg A, Skjaerven R, Forman J, Gissler M, Nygren KG, Tiitinen A. Perinatal outcomes of children born after frozen-thawed embryo transfer: a Nordic cohort study from the CoNARTaS group. *Hum Reprod* 2013;**28**:2545–2553.
- White CR, Denomme MM, Tekpetey FR, Feyles V, Power SG, Mann MR. High frequency of imprinted methylation errors in human preimplantation embryos. *Sci Rep* 2015;**5**:17311.
- Wise LA, Mikkelsen EM, Sørensen HT, Rothman KJ, Hahn KA, Riis AH, Hatch EE. Prospective study of time to pregnancy and adverse birth outcomes. *Fertil Steril* 2015;**103**:1065–1073.
- Xu J, Sinclair KD. One-carbon metabolism and epigenetic regulation of embryo development. *Reprod Fertil Dev* 2015. doi:10.1071/RD14377.
- Yeung EH, Druschel C. Cardiometabolic health of children conceived by assisted reproductive technologies. *Fertil Steril* 2013;**99**:318–326.
- Zandstra H, Van Montfoort AP, Dumoulin JC. Does the type of culture medium used influence birthweight of children born after IVF? *Hum Reprod* 2015;**30**:530–542.
- Zhang Y, Cui Y, Zhou Z, Sha J, Li Y, Liu J. Altered global gene expressions of human placentae subjected to assisted reproductive technology treatments. *Placenta* 2010;**31**:251–258.
- Zhou J, Liu H, Gu H-T, Cui Y-G, Zhao N-N, Chen J, Gao L, Zhang Y, Liu J-Y. Association of cardiac development with assisted reproductive technology in childhood: a prospective single-blind pilot study. *Cell Physiol Biochem* 2014;**34**:988–1000.